



Evidence-Based Care Guideline

For management of
Acute Bacterial Sinusitis
in children 1 to 18 years of age¹

Original Publication Date: 04-27-01

Revision Publication Date: 7-7-06

Target Population

Inclusions: These guidelines are intended primarily for use in children 1 to 18 years of age with suspected acute bacterial sinusitis (ABS).

Exclusions: These guidelines do not address all considerations needed to manage children:

- under 1 year of age
- with chronic sinusitis
- with identified or suspected periorbital, orbital or intracranial abscess
- with cystic fibrosis
- with underlying anatomic paranasal abnormalities
- with ciliary dyskinesia
- with immune deficiencies

Target Users

Includes but is not limited to (in alphabetical order):

- allergists
- clinicians caring for inpatients
- emergency medicine physicians
- ophthalmologists²
- otolaryngologists³

¹ Please cite as: **Acute Bacterial Sinusitis Guideline Team, Cincinnati Children's Hospital Medical Center:** Evidence-based care guideline for medical management of Acute Bacterial Sinusitis in children 1 through 17 years of age, <http://www.cincinnatichildrens.org/svc/alpha/h/health-policy/ev-based/sinus.htm>, Guideline 16, pages 1-17, July 7, 2006.

² ophthalmologist = eye physician

³ otolaryngologist = ENT, ear/nose/throat physician

- patient care staff
 - nurse practitioners
 - nurses
 - pharmacists
- patients and families
- physicians in training
- primary care providers
- radiologists

Introduction

References in parentheses () Evidence strengths in [] (See last page for definitions)

Little new evidence on the management of acute bacterial sinusitis (ABS) in children has been published since the original publication of this guideline. Changes for this revision include:

- the guideline recommendations have been updated to reflect current antibiotic prescribing patterns
- stronger evidence is cited to support recommendations regarding:
 - the lack of benefit of over-the-counter cough preparations, and
 - the need to discuss parental and physician views and expectations for the office visit in the management of ABS
- a new appendix listing likelihood ratios (LRs) for clinical signs and symptoms and diagnostic studies for ABS in children, and an introduction to the use of these LRs.

ABS is a bacterial infection of the paranasal sinuses lasting less than 30 days in which symptoms resolve completely (*Wasserfallen 2004 [S], AAP 2001 [S]*). It is a disorder that frequently presents in the primary care setting (*AAP 2001 [S], Brook 2000 [S,E], Dowell 1998a [E]*). See Table 1 for clinical signs and symptoms consistent with a diagnosis of ABS.

Sinusitis accounted for 9% of all pediatric antibiotic prescriptions written in 2002. Studies of the epidemiology and natural history of the condition suggest that spontaneous resolution of ABS ranges from 30% to 80% (*Anon 2004 [S,E]*).

A variety of conditions affecting the upper respiratory tract can confound the accurate diagnosis and appropriate treatment of ABS. An accurate differentiation of the more common upper respiratory infections (URIs) from sinusitis will reduce inappropriate use of antibiotics in children with URIs

Table 1: Clinical Signs and Symptoms Consistent With a Diagnosis of Acute Bacterial Sinusitis

Acute bacterial sinusitis	Acute severe bacterial sinusitis										
<p>Characterized by:</p> <ul style="list-style-type: none"> ● persistence of upper respiratory symptoms for greater than 10 days without improvement (<i>Wald 1984 [B], Wald 1981 [B], Brook 2000 [S,E], Isaacson 1996 [S], Wald 1994 [S]</i>) <p>AND</p> <ul style="list-style-type: none"> ● nasal congestion and nasal discharge of any quality (i.e., may be either thin and milky or thick and purulent) (<i>Wald 1984 [B], Wald 1981 [B], Aitken 1998 [C], Kogutt 1973 [D], McLean 1970 [D], Gungor 1997 [S], Wald 1994 [S], Williams 1993 [S], Fireman 1992 [S], Ott 1991 [S], Dowell 1998a [E]</i>) ● persistent cough which is often more severe at night is often present (<i>Wald 1984 [B], Wald 1994 [S]</i>) <p>Note: Less common complaints may include:</p> <ul style="list-style-type: none"> ● low-grade fever (<i>Wald 1984 [B], Aitken 1998 [C], McLean 1970 [D]</i>) ● sore throat or ear discomfort (<i>Brook 2000 [S,E], Zacharisen 1998 [S]</i>) ● fatigue (<i>Brook 2000 [S,E], Zacharisen 1998 [S]</i>) ● malodorous breath (<i>Wald 1991 [C], Lusk 1997 [S]</i>) ● intermittent periorbital edema or facial swelling (<i>Zacharisen 1998 [S], Wald 1994 [S]</i>) ● facial or tooth pain (<i>Wald 1991 [C], Zacharisen 1998 [S], Isaacson 1996 [S], Wald 1994 [S]</i>) 	<p>A less common presentation and may present with the same symptoms described to the left but is differentiated from ABS by severity, rather than persistence, of symptoms:</p> <ul style="list-style-type: none"> ● ill-appearing child (<i>AAP 2001 [S], Fireman 1992 [S]</i>) <p>AND</p> <ul style="list-style-type: none"> ● fever higher than 39° C (102.2° F) (<i>Wald 1984 [B], AAP 2001 [S], Wald 1994 [S], Fireman 1992 [S]</i>) <p>AND</p> <ul style="list-style-type: none"> ● purulent nasal discharge, typically 3 to 4 days duration (<i>AAP 2001 [S], Wald 1994 [S], Fireman 1992 [S]</i>) 										
<p>Other conditions that present with symptoms similar to ABS to be considered in the differential diagnosis include:</p> <table style="width: 100%; border: none;"> <tr> <td>recurrent viral URI</td> <td>allergic rhinitis</td> <td>cough-variant asthma</td> <td>enlarged adenoids</td> </tr> <tr> <td>deviated nasal septum</td> <td>choanal atresia</td> <td>nasal foreign body</td> <td>neoplasm</td> </tr> </table> <p>(<i>Williams 1993 [S]</i>)</p> <table style="width: 100%; border: none;"> <tr> <td>gastroesophageal reflux disease</td> <td>nasal polyp</td> </tr> </table> <p>(<i>Isaacson 1996 [S], Williams 1993 [S]</i>)</p>		recurrent viral URI	allergic rhinitis	cough-variant asthma	enlarged adenoids	deviated nasal septum	choanal atresia	nasal foreign body	neoplasm	gastroesophageal reflux disease	nasal polyp
recurrent viral URI	allergic rhinitis	cough-variant asthma	enlarged adenoids								
deviated nasal septum	choanal atresia	nasal foreign body	neoplasm								
gastroesophageal reflux disease	nasal polyp										

(*Anon 2004 [S,E], Dowell 1998b [E], Dowell 1998a [E]*). See Table 1. Risk of serious bacterial complications must also be considered (*Oxford 2005 [D]*).

The amount of quality evidence for diagnosis and treatment of pediatric ABS is limited compared to the frequency of its occurrence (*Nyquist 1998 [D], Wasserfallen 2004 [S], Contopoulos-Ioannidis 2003 [S], Lau 1999 [S]*). As the causative organisms in pediatric ABS and otitis media are identical, where evidence was minimal or non-existent, literature from pediatric otitis media studies was extrapolated for use in treatment recommendations (*Parsons 1996 [E]*). In the absence of quality evidence, expert local consensus was used. Because scant evidence exists concerning the management of a child less than 1 year of age with ABS, this guideline is limited to children 1 year of age and older.

Areas of uncertainty offering challenges in the management of ABS include:

- clinical criteria for defining ABS in children;
- specific parameters for effective antibiotic management of ABS in children.

The objectives of this guideline are to:

- improve the recognition of clinical signs and symptoms consistent with the diagnosis of ABS
- improve the use of appropriate radiologic studies in the diagnosis of ABS
- improve the judicious use of antibiotics in the treatment of ABS
- outline parameters for appropriate referral to and integration of subspecialty services.

Etiology

Acute bacterial sinusitis is an inflammation of the paranasal sinus mucous membrane caused by bacterial overgrowth in a closed cavity. The pathogens most commonly isolated in studies that have included maxillary sinus aspirate in children are listed in Table 2. Due to various parameters which affect prevalence, current surveillance monitoring in the local community might demonstrate a different distribution.

Table 2: Pathogens in Pediatric Acute Bacterial Sinusitis

Organism	Proportion (%)
<i>Streptococcus pneumoniae</i>	25 to 30%
<i>Haemophilus influenzae</i>	15 to 20%
<i>Moraxella catarrhalis</i>	15 to 20%
<i>Streptococcus pyogenes</i>	Up to 5%
No identified organism	30%

(Wald 1984 [B], Wald 1981 [B])

Note: Pathogens may be different in individuals with chronic sinusitis and in those with underlying chronic conditions such as cystic fibrosis (Brook 1981 [C], Muntz 1991 [D]).

Incidence

Children two to five years of age average six to eight URIs per year. Up to 7% of these pediatric URIs become complicated by ABS (Wald 1991 [C]). It is estimated that antibiotic treatment exclusively for sinusitis accounts for 5% of office visits by preschool children seen during winter months (Aitken 1998 [C]).

Local experience in one practice of 20 pediatricians indicates that 10% of total office visits seen in a single year were related to sinusitis (acute, chronic or follow-up) (Group Health Associates 2004 [O]).

Predisposing Factors:

The most common predisposing factors are viral URIs, allergic inflammation (Wald 1981 [B]), and exposure to smoke (Kakish 2000 [C]).

Guideline Recommendations

Assessment and Diagnosis

See Table 1 for clinical signs and symptoms consistent with a diagnosis of ABS.

Clinical Assessment

- It is recommended that the diagnosis of ABS be made clinically in the presence of a constellation of signs and symptoms of **at least 10 days duration without improvement** (Wald 1984 [B], Wald 1981 [B], Aitken 1998 [C], Wald 1991 [C]). No single symptom or sign is specific for the diagnosis of ABS. See Appendix 1 for likelihood ratios for clinical signs and symptoms.

Note 1: The 10-day duration is suggested because it has been shown that, in most children with uncomplicated URI, improvement is seen on average by 10 days (Wald 1984 [B], Wald 1981 [B], Wald 1991 [C]).

Note 2: A less common presentation, acute severe bacterial sinusitis, represents a more

toxic form of ABS in which severity of symptoms, rather than persistence of symptoms, is consistent with the diagnosis (Wald 1994 [S], Fireman 1992 [S]). See Table 1.

- It is recommended that the character of the nasal discharge **not** be used to make a diagnosis or as an indication for antibiotic treatment. The quantity, quality, and color of nasal discharge are not helpful in differentiating ABS from other upper respiratory illnesses (e.g. common cold, allergic rhinitis) (Wald 1981 [B], Aitken 1998 [C], McLean 1970 [D], Gungor 1997 [S], Wald 1994 [S]).

Note: Physical exam is likely to reveal purulent nasal discharge and/or posterior oropharyngeal drainage. These findings, however, are non-specific and of little diagnostic usefulness (Wald 1981 [B], McLean 1970 [D], Williams 1993 [S], Fireman 1992 [S]).

Radiologic Assessment

- It is recommended that radiologic studies **not** be routinely obtained in the initial management of patients with suspected uncomplicated ABS (Engels 2000 [M], Schwartz 2001 [C], AAP 2001 [S], McAlister 2000 [E], Diament 1992 [E]). See Appendix 1 for likelihood ratios for radiologic studies.

Note 1: Abnormalities of the paranasal sinuses are found frequently on conventional radiographs and computed tomography (CT) scans in children **without** clinical evidence of sinusitis (see Table 3) (Rak 1991 [C], Glasier 1989 [C], Diament 1987 [C], Glasier 1986 [C], Odita 1986 [C], Shopfner 1973 [C], Maresh 1940 [D]).

Note 2: The presence of a URI alone (without sinusitis) can result in mucosal thickening and abnormal findings in the paranasal sinuses on plain radiographs and CT scans (Glasier 1989 [C], Glasier 1986 [C], Shopfner 1973 [C], Gwaltney 1994 [D]).

Note 3: Imaging findings may persist well after symptoms improve. CT abnormalities with the common cold may last up to two weeks after symptomatic improvement (Gwaltney 1994 [D]). Magnetic resonance imaging (MRI) changes in patients with symptoms of ABS may last more than eight weeks (Leopold 1994 [C]).

Note 4: "Limited" sinus CT lacks sensitivity in identifying air-fluid levels (Gross 1991 [C]), suboptimally visualizes the osteomeatal complex 30% of the time, and misses 20 to 30% of the findings found on full CT (Wippold 1995 [C]).

4. It is recommended, for older children with persistent clinical findings after unsuccessful therapy, or for children with clinical evidence of orbital or intracranial complications of ABS, that the decision to perform radiologic studies be made in collaboration with the consulting ophthalmologist or otolaryngologist (*Oxford 2005 [D]*, *Vazquez 2004 [D]*, *AAP 2001 [S]*, *Local Expert Consensus [E]*). See Table 4 for radiologic modalities and Table 5 for description of complications.

Note 1: An otolaryngology or ophthalmology consultation prior to obtaining radiologic studies in this patient population may reduce the need for an early study and limit repeat radiation exposure (*Local Expert Consensus [E]*).

Note 2: A clear or normal Water's view (occipitontal) may be helpful in ruling out significant maxillary sinus disease (*Ros 1995 [D]*, *Lau 1999 [S]*, *Wald 1988 [E]*).

Table 3: Abnormal Imaging in Children without Upper Respiratory Symptoms

Age Range	Imaging Mechanism	% Abnormal
*6 mo to 15yrs	Plain films	15 to 57%
**Infants and children	CT scan	18 to 67%
***15 to 85yrs	MRI	80%

*(*Diament 1987 [C]*, *Odita 1986 [C]*, *Shopfner 1973 [C]*, *Maresh 1940 [D]*)

**(*Glasier 1989 [C]*, *Glasier 1986 [C]*)

***(*Rak 1991 [C]*)

Table 4: Radiologic Modalities for Suspected Complications of Acute Bacterial Sinusitis

Indication	Modality
Suspected subperiosteal or orbital abscess	Contrast enhanced CT scan of orbits (thin section)
Suspected intracranial complications	Contrast enhanced CT or MRI of brain

(*Vazquez 2004 [D]*, *AAP 2001 [S]*, *McAlister 2000 [E]*, *Local Expert Consensus [E]*)

Laboratory Assessment

5. It is recommended that routine laboratory testing such as a complete blood count (CBC) or nasopharyngeal culture **not** be obtained in the initial evaluation in children with uncomplicated ABS (*Clement 1998 [E]*). See Appendix 1 for likelihood ratios for laboratory studies.

Note: Organisms recovered from nasopharyngeal washings and throat culture do not reflect the organisms found in sinus aspirate (*Wald 1981 [B]*).

6. It is recommended that sinus aspiration and bacterial culture **not** be obtained for use in the initial evaluation and management of the child with uncomplicated ABS. They are recognized as the "gold standard" for definitive diagnosis of bacterial sinusitis and may need to be considered under the following situations (*Wald 1981 [B]*):
- severe illness or toxic-looking child
 - immunocompromised child
 - presence of suppurative or intracranial complications.

Management

General

The treatment of pediatric ABS is best considered in light of the duration and severity of symptoms and the increasing prevalence of resistant strains of a common sinus pathogen, *Streptococcus pneumoniae*. The treatment recommendations for this guideline were developed with a focus on antimicrobial activity against *S. pneumoniae* in an era of increasing penicillin resistance. It is prudent for clinicians to consider use of the most narrow-spectrum agent that is active against the likely pathogens for the initial antimicrobial treatment of ABS in children (*Dowell 1998a [E]*).

See Appendix 2 for antibiotic dosages.

Antibiotic Treatment

7. It is recommended that high-dose amoxicillin (80 to 90 mg / kg / day) or amoxicillin-clavulanate (with high-dose amoxicillin component) be first-line therapy for most patients with pediatric ABS (*Wald 1986 [B]*, *AAP 2001 [S]*, *Nash 2001 [S]*, *Dowell 1999 [E]*, *Friedland 1994 [E]*, *Local Expert Consensus [E]*). Treatment duration is 10 to 14 days to minimize the development of bacterial resistance (*Morris 2005 [M]*, *Local Expert Consensus [E]*). See Appendix 2.

Note 1: Approximately 65% of the *S. pneumoniae* isolated from non-sterile sites of children in Cincinnati in outpatient settings are resistant to penicillin (*Cincinnati Children's Hospital Medical Center 2005 [O]*).

Note 2: It is recognized that the rates of *S. pneumoniae* resistance to penicillin are increasing nationally and locally (*Butler 1996 [C]*, *Breiman 1994 [C]*) and failure with amoxicillin is likely to be due to resistant *S. pneumoniae*, *H. influenzae*, or *M. catarrhalis* (*Whitney 2000 [D]*). Resistance of *S. pneumoniae* to penicillin (including amoxicillin) is mediated through alterations in the penicillin-binding proteins. Using high doses of amoxicillin saturates the

penicillin-binding proteins, and is therefore considered a reasonable antibiotic option (Schrag 2001 [A], Dagan 2001 [C], Dowell 1999 [E]). The clavulanic acid component of amoxicillin-clavulanate is active against resistant *H. influenzae* and *M. catarrhalis* (β -lactamase enzyme) (Dagan 2000 [A], Wald 1986 [B]).

Note 3: Toxic-appearing children who demonstrate poor tolerance of oral intake may require initial parenteral therapy either as an outpatient or a short inpatient stay. Reassessment after initial stabilization may avoid unnecessary imaging and referral early in the course of therapy (Local Expert Consensus [E]).

8. It is recommended that cefuroxime, cefpodoxime, and cefdinir be second-line therapy for pediatric ABS (Pichichero 1997 [C], Felmingham 2005 [D], Jacobs 2003 [D], Jacobs 1999 [D], Anon 2004 [S,E], Dowell 1999 [E]). Treatment duration is 10 to 14 days to minimize the development of bacterial resistance (Morris 2005 [M], Local Expert Consensus [E]). See Appendix 2.
9. It is recommended, if clinical failure with a second-line agent occurs, that alternative agents or combination therapy be considered:
 - IM ceftriaxone (5 days)
 - combination therapy with adequate gram-positive and -negative coverage, such as clindamycin plus cefixime
 (Anon 2004 [S,E], AAP 2001 [S], Local Expert Consensus [E]). See Appendix 2.
10. It is recommended, in the penicillin-allergic patient, that the following be used:
 - non-type I^d: cefdinir, cefuroxime, or cefpodoxime
 - type I^e: clarithromycin or azithromycin
 (AAP 2001 [S], Local Expert Consensus [E]). See Appendix 2.

Note: Macrolides, azilides, and sulfa containing agents are not considered standard therapeutic agents due to either a lack of efficacy data, increasingly resistant *S. pneumoniae* or both (Dagan 2000 [A], Nelson 1994 [C], Wu 2004 [D], Gay 2000 [D], AAP 2001 [S]).

^d Non-type I penicillin allergy: more common; characterized by symptoms such as maculopapular, polymorphous rash, arthralgia, or emesis.

^e Type I penicillin allergy: IgE-mediated; rare; anaphylactic reactions result in urticaria, pruritis, laryngeal edema, bronchospasm, cardiovascular collapse, and, potentially, death.

Symptomatic Treatment

11. It is recommended that common agents for symptomatic treatment of cough or congestion (i.e. reduction in frequency or severity), **not** be used in the routine management of patients with ABS (Schroeder 2004 [M], Bernard 1999 [B], Davies 1999 [B], Chang 1998 [B], McCormick 1996 [B], Taylor 1993 [B], Gadomski 1992 [O], Local Expert Consensus [E]).

Note 1: Studies measuring a decrease in frequency, severity, and time to resolution of cough or congestion in children with symptoms from URI found no significant difference between any of the therapeutic interventions and placebo. The therapies evaluated were antitussives, mucolytics, inhaled steroids, inhaled and oral beta₂-agonists, antihistamines / decongestants (brompheniramine, phenylephrine, phenylpropanolamine, dextromethorphan/ guaifenesin, oxymetolazine or “afrin”), and morphine derivatives (codeine) (Schroeder 2004 [M], Paul 2004 [A], Bernard 1999 [B], Davies 1999 [B], Chang 1998 [B], McCormick 1996 [B], Taylor 1993 [B], Gadomski 1992 [O]).

Note 2: One previously common ingredient (phenylpropanolamine) of symptomatic treatment preparations has been associated with stroke, and most antihistamines, decongestants, and antitussives have not been FDA approved in children (Kernan 2000 [D], AAP 1997 [S,E]).

Note 3: Although hypertonic and normal saline and balanced physiological saline nasal washes are commonly used in postoperative patients and in children with chronic sinusitis (Shoseyov 1998 [B], Pigret 1996 [B], Nuutinen 1986 [C]) there is no evidence for their effectiveness in pediatric ABS.

Follow up

12. It is recommended that follow-up assessment for expected clinical response occur by 72 hours of antimicrobial therapy. A lack of expected clinical improvement may indicate that a change of antibiotic is necessary (Wald 1986 [B], Dowell 1999 [E], Local Expert Consensus [E]).

Consults and Referrals

Although children with the complications discussed below (see Table 5) are listed as exclusions to this guideline, recommendations are included here to assist the practitioner in decisions regarding consultation to specialists for these key complications.

13. It is recommended that an otolaryngology and/or ophthalmology consultation be sought when signs of impending suppurative complications of ABS are present (AAP 2001 [S], Local Expert Consensus [E]). Such complications are rare but very serious and often result from orbital or intracranial spread of infection (Oxford 2005 [D], Rosenfeld 1994 [D]).

Note 1: Preseptal cellulitis, involving only tissue anterior to the orbital septum, manifests as lid edema/erythema, conjunctivitis, and fever. It may be treated with oral antibiotics and close follow up except where toxicity or specific symptoms preclude adequate antimicrobial effectiveness by mouth (AAP 2001 [S]).

Note 2: Consultation prior to imaging limits repeat radiation exposure (Local Expert Consensus [E]).

14. It is recommended that otolaryngology consultation be considered in cases of a moderately to severely ill child with suspected acute frontal or sphenoid sinusitis because of the potential for intracranial spread. Infection arising in either site will generally occur in a relatively older age group (>6 years), and based on the developmental anatomy of these sinuses, the clinical presentation is likely to be more severe (Oxford 2005 [D], Herrmann 2004 [D], Wolf 1993 [F]).

Note 1: Acute frontal sinusitis manifests as an intense frontal headache with tenderness over the sinus itself. Spread of infection anteriorly produces periosteal edema and osteomyelitis and may manifest as doughiness of the forehead skin, known as Pott’s puffy tumor. Spread of infection to the cranial vault results in meningitis or intracranial abscess (Oxford 2005 [D]).

Note 2: Acute isolated sphenoid sinusitis is rare, with an estimated incidence of <1% of all sinusitis cases (Hnatuk 1994 [S], Fearon 1979 [S], Wyllie 1973 [S]). Acute sphenoid sinusitis represents an elusive diagnosis (Myer 1982 [S], Sellars 1975 [S], Postma 1995 [E]), as signs and symptoms are more variable and non-specific than those of frontal sinus disease. Nasal symptoms may be absent. Headache is severe, deep-seated and worse at night, with the pain radiating to any craniofacial region (Myer 1982 [S], Sellars 1975 [S]). Suppurative complications may involve any of the vital juxtaposing structures, including the cavernous sinus, intracranial cavity, orbit, pituitary gland, or abducens nerve.

Table 5: Complications of Pediatric Acute Bacterial Sinusitis

Complication	Signs and Symptoms	Intervention
Orbital cellulitis	Fever, lid edema/erythema, conjunctivitis, chemosis, altered acuity, proptosis, ophthalmoplegia, pain with eye movement, tenderness to palpation	<u>IV antibiotics</u> <u>Consult:</u> • otolaryngology and/or ophthalmology
Subperiosteal abscess	Above, with proptosis and ophthalmoplegia prominent features; +/- globe displacement laterally or superiorly	<u>Imaging:</u> • decision to image made in collaboration with consulting specialist
Orbital abscess	Same as for orbital cellulitis, with proptosis and chemosis prominent features; severe impairment of vision	
Cavernous sinus thrombosis and/or Intracranial infection	Spiking fevers, cranial neuropathy, mental status changes	In addition to above: <u>Consult:</u> • neurosurgery • infectious diseases

(Oxford 2005 [D], Vazquez 2004 [D], AAP 2001 [S])

Parental Expectations and Education

15. It is recommended that, for a child with ABS, physicians explore parental expectations concerning the office visit, parental knowledge regarding respiratory infections, and preventive behavior (Mangione-Smith 1999 [C], Barden 1998 [C], Macfarlane 1997 [C], Varonen 2004 [O], Local Expert Consensus [E]). Topics for discussion may include:


- the natural history of URIs / ABS (Roberts 1983 [A], Hamm 1996 [C])
- diagnostic uncertainty (Varonen 2004 [O])
- viral and bacterial sources of ABS
- role of antibiotics (Hamm 1996 [C])
 - appropriate use of antibiotics (Mangione-Smith 1999 [C], Barden 1998 [C], Macfarlane 1997 [C])
 - persistent or severe infections (Garbutt 2001 [B], Wald 1986 [B])
- bacterial resistance (Trepka 2001 [C])
- lack of proven efficacy for over-the-counter medications for symptom relief (Schroeder 2004 [M])
 - managing cough symptoms
- observation for complications of ABS
- prevention of URIs may decrease risk of ABS
 - handwashing (Morton 2004 [A], Roberts 2000 [A])
 - annual influenza vaccination (Loughlin 2003 [D]).

Future Research Agenda

It has been noted in a recent meta-analysis that there is a paucity of high quality primary data in the area of diagnosis and management of children suspected to have ABS (*Ioannidis 2001 [M]*). Following are some clinical questions related to guideline recommendations and of potential interest to CCHMC investigators:

1. In children with suspected ABS, what clinical signs and symptoms are most useful in identifying children most likely to benefit from antibiotics?
2. In children with suspected ABS < 10 days duration, what clinical signs and symptoms are most useful in the diagnosis of acute severe bacterial sinusitis?
3. In children with ABS, what criteria define judicious use of antibiotics, given high spontaneous cure rates for ABS and increasing global antibiotic resistance?
4. In the local community, what is the relative prevalence of *S. pneumoniae* and *H. influenzae* in children with ABS, and what is the antibiotic resistance profile of these pathogens?
5. How do clinical signs and symptoms and radiologic imaging correlate with bacterial and viral cultures from sinus puncture?

Treatment Algorithm for Acute Bacterial Sinusitis (ABS)



Appendix 1: Likelihood Ratios (LR)

A. Definition

When a health care provider evaluates a patient, he/she determines their own “best guess” for how likely a disease is present (or not present) at that time.

This “best guess” is dependent on:

- the disease prevalence in the community,
- the patient’s underlying medical status and current presentation, and
- the health care provider’s experience and knowledge of the literature.

This best guess is actually the **pre-test probability**.

What health care providers are looking for is a test which will increase (or decrease) the **likelihood** of disease in that patient, thus allowing them to decide to treat, not treat, or pursue further diagnostic work up.

This change in “best guess” after diagnostic testing results in the **post-test probability** of disease.

Finding the post-test probability is done easily with **likelihood ratios** (see next page) and a **nomogram** (see figure).

What the likelihood ratio nomogram actually does is change pre-test probability to pre-test odds, multiply by the LR to get post-test odds, then change post-test odds back to post-test probability...all without having to manually perform complex calculations.

Fortunately, likelihood ratios (**LRs**) are determined by inherent properties of the diagnostic test, not the prevalence of disease in the population. By knowing the likelihood ratio(s) for any or all given tests, health care providers can determine which test is most informative and appropriate to use.

B. Rule of thumb:

An **LR value**

- greater than 10 is very helpful in increasing diagnostic certainty
the presence of clinical sign is 10 times more likely to be present in a child with ABS than in a child without ABS
- of 1 is not helpful
the presence of clinical sign is just as likely to be present in a child with ABS as in a child without ABS
- less than 0.2 is very helpful in ruling out the condition
the presence of clinical sign is one-fifth as likely to be present in a child with ABS as in a child without ABS

C. How to calculate a Likelihood Ratio from sensitivity and specificity data:

LR for a positive result = (sensitivity) / (1-specificity)

LR for a negative result = (1-sensitivity) / (specificity)

For more information on LRs see: <http://www.cebm.utoronto.ca/glossary/lrs.htm#top>

See likelihood ratios for common signs and symptoms for use in diagnosing acute bacterial sinusitis next page.




Figure: Likelihood Ratio Nomogram

Appendix 1: Likelihood Ratios (LR), continued**D. Likelihood ratios (LRs) for clinical signs/symptoms of acute bacterial sinusitis in children**

Sign/Symptom	Age range	LR (+)
Presence of symptoms for at least 10 days AND At least one of the following present: <ul style="list-style-type: none"> • blocked or stuffy nose • headaches or face pain • coughing during the day • coughing during the night • yellow or green mucus from the nose 	6 months to 18 years	11.6

(Garbutt 1999 [C])

E. Likelihood ratios (LRs) for diagnostic study results of acute bacterial sinusitis in children

Imaging or Aspiration Test	“Gold standard” comparison	LR (+)	LR (-)
CT scan of ethmoid sinuses in infants: opacification* (Ioannidis 2001 [M], Glasier 1989 [C])	Clinically diagnosed URI	1.7	0.5
CT scan of maxillary sinuses in infants: opacification* (Ioannidis 2001 [M], Glasier 1989 [C])	Clinically diagnosed URI	1.5	0.3
Plain radiograph: any abnormality** (Ioannidis 2001 [M], Jannert 1982 [C])	Specified clinical criteria	2.8	0.3
Plain radiograph: any abnormality** (Ioannidis 2001 [M], Van Buchem 1992 [C])	Non-clear fluid in sinus aspirate	1.1	0.9
Plain radiograph: any abnormality** (Ioannidis 2001 [M], Watt-Boolsen 1977 [C])	Any fluid in sinus aspirate	2.7	0.2
Sinus aspirate: non-clear fluid (Ioannidis 2001 [M], Van Buchem 1992 [C])	Pathogens in sinus aspirate	0.9	1.0
Ultrasound: any abnormality (Ioannidis 2001 [M], Van Buchem 1992 [C])	Any abnormality** on plain radiograph	1.7	0.9
Ultrasound: any abnormality (Ioannidis 2001 [M], Van Buchem 1992 [C])	Non-clear fluid in sinus aspirate	0.5	1.2
Radiograph with sinus fluid or opacity (Engels 2000 [M])	Sinus puncture	3.65	0.34
Radiograph with sinus fluid or opacity or mucous membrane thickening (Engels 2000 [M])	Sinus puncture	2.31	0.16

*excluding hypoplasia

**typically including mucous thickening, opacification, or air-fluid level

NOTE: The figures in these tables highlight the fact that the presence of clinical signs and symptoms of acute bacterial sinusitis (ABS) are generally more clinically useful in making the diagnosis of ABS than radiologic or laboratory studies. The clinical signs/symptoms positive likelihood ratio of 11.6 indicates a much greater probability of ABS when clinical findings are present, than the positive likelihood ratios of radiologic or laboratory studies, which range from 0.5 to 3.65. Also, a negative radiologic or laboratory test result is not helpful in ruling out ABS, as these negative likelihood ratios range from 0.16 to 1.2.

Rule of thumb:**An LR value**

- greater than 10 is very helpful in increasing diagnostic certainty
- of 1 is not helpful
- less than 0.2 is very helpful in ruling out the condition

Appendix 2: Antibiotic Therapy for Acute Bacterial Sinusitis

Antibiotic	Dose, Frequency & Max Daily Dose	Oral Dosage Forms	Relative Cost	Comments
First-line therapy				
amoxicillin	80 to 90 mg / kg / day Max daily dose 2 gm taken as: 40 to 45 mg / kg BID or 25 to 30 mg / kg TID	Suspension (per 5mL): 125, 200, 250 or 400 mg Capsules 250 or 500 mg Chewable tablet: 125, 200, 250, 400 mg Tablet: 500 or 875 mg	Low	<ul style="list-style-type: none"> High-dose for resistant <i>S. pneumoniae</i> 400 mg / 5ml formulation not covered by Ohio Medicaid (June, 2006)
amoxicillin-clavulanate (Augmentin®)	(amoxicillin) 80 to 90 mg / kg / day Max daily dose: 2 gm taken as: 40 to 45 mg / kg BID	amoxicillin component: Suspension (per 5mL) 125, 200, 250, 400, 600 mg Chewable tablet: 125, 200, 250, 400 mg Tablet: 500, 875 mg	High	Maximum dose of clavulanate not to exceed 6.4 mg / kg / day, to minimize diarrhea
Second-line therapy (first-line for non-type I^f penicillin-allergic patient)				
cefuroxime (Ceftin®)	NOTE: Tablets and suspension are not bioequivalent and are not substitutable on a mg-for-mg basis.		High	Unpleasant taste (<i>Steele 1997 [O]</i>)
	Suspension: 30 mg / kg / day Max daily dose: 1 gm taken as: 15 mg / kg BID Dosage form: (per 5 mL): 125 or 250 mg	Tablet: 500 mg / day taken as: 250 mg BID Dosage form: 250, 500 mg		
cefepodoxime (Vantin®)	10 mg / kg / day Max daily dose: 800 mg taken as: 5 mg / kg BID	Suspension (per 5mL): 50, 100 mg Tablet: 100, 200 mg	High	Not covered by Ohio Medicaid (June, 2006)
cefdinir (Omnicef®)	14 mg / kg / day Max daily dose: 600 mg taken as: 7 mg / kg BID or 14 mg / kg once a day	Suspension (per 5 mL): 125, 250 mg Capsule: 300 mg	High	
Alternative Agents				
clindamycin (Cleocin®)	30 mg / kg / day Max daily dose: 1.8 g taken as: 10 mg / kg TID	Suspension (per 5 mL): 75 mg Capsule: 75, 150 or 300 mg	Med	<ul style="list-style-type: none"> If <i>S. pneumoniae</i> is identified as a pathogen (<i>Anon 2004 [S,E], AAP 2001 [S]</i>) Use in combination with cefixime or other gram-negative coverage (<i>Anon 2004 [S,E]</i>)
ceftriaxone (Rocephin®)	50 mg / kg Max daily dose: 1 gm taken as: once daily for 5 days (<i>Anon 2004 [S,E]</i>)	Intramuscular	High	For treatment failure (<i>Anon 2004 [S,E], AAP 2001 [S]</i>)
Type I^g penicillin-allergic patient therapy				
clarithromycin (Biaxin®)	15 mg / kg / day Max daily dose: 1 gm taken as: 7.5 mg / kg BID	Suspension (per 5 mL): 125 or 250 mg 250 and 500 mg tablets	High	<ul style="list-style-type: none"> Not covered by Ohio Medicaid (June, 2006) Unpleasant taste (<i>Steele 1997 [O]</i>)
azithromycin (Zithromax®)	day 1 = 10 mg / kg day 2 to 5 = 5 mg / kg Max daily dose: 500 mg taken: once a day or 20 mg / kg once daily for 3 days Max daily dose: 500 mg	Suspension (per 5 mL): 100 or 200 mg Tablet 250, 500 mg	High	<ul style="list-style-type: none"> Not covered by Kentucky Medicaid (June, 2006) Not used as standard therapeutic agents due to lack of efficacy and/or increasingly resistant <i>S. pneumoniae</i> (<i>Gay 2000 [D]</i>)

^f Non-type I penicillin allergy: more common; characterized by symptoms such as maculopapular, polymorphous rash, arthralgia, or emesis.

^g Type I penicillin allergy: IgE-mediated; rare; anaphylactic reactions result in urticaria, pruritis, laryngeal edema, bronchospasm, cardiovascular collapse, and, potentially, death.

Acute Bacterial Sinusitis Team Members 2005-6

Community Physician

*Stephen Bird, MD, Chair, Pediatrics

CCHMC Physicians and Practitioners

*Amal Assa'ad, MD, Allergy/Immunology

Michael Gerber, MD, Infectious Diseases

*Bernadette Koch, MD, Radiology

*Paul Willging, MD, Otolaryngology

*Constance West, MD, Ophthalmology

Tiffany Raynor, MD, Otolaryngology

Patrick Brady, MD, Resident

Patient Services

*Dawn Butler, PharmD, Pharmacy

*Donna Hillman, RN, Emergency Dept

Division of Health Policy Clinical Effectiveness Support

Eloise Clark, MPH, Facilitator

*Kieran Phelan, MD, Methodologist, General Pediatrics

Edward Donovan, MD, Medical Director, Clinical Effectiveness

Eduardo Mendez, RN, MPH, Dir. Evidence-Based Care

Danette Lopp-Stanko, MA, MPH, Epidemiologist

Carol Tierney, RN, MSN, Education Specialist

*Kate Rich, Lead Decision Support Analyst

Deborah Hacker, RN, Medical Reviewer

All Team Members and Clinical Effectiveness support staff listed above have signed a conflict of interest declaration.

Ad hoc Advisors

*Uma Kotagal, MBBS, MSc, VP, Division Director

*Alan Brody, MD, Radiology

*Thomas DeWitt, MD, General & Community Pediatrics

*Richard Ruddy, MD, Emergency Medicine

*Michael Farrell, MD, Chief of Staff

*Beverly Connelly, MD, Director Infectious Diseases

Cheryl Hoying, RN, PhD, Sr. VP, Patient Services

*Barbarie Hill, Pratt Library

*Member of previous Acute Bacterial Sinusitis guideline development Team

Development Process

The process by which this guideline was developed is documented in the [Guideline Development Process Manual](#); a Team Binder maintains minutes and other relevant development materials. The recommendations contained in this guideline were formulated by an interdisciplinary working group which performed systematic and critical literature reviews, using the grading scale that follows, and examined current local clinical practices.

To select evidence for critical appraisal by the group, the Medline, EmBase and the Cochrane databases were searched for dates of Jan 2000 through December, 2005 to generate an unrefined, "combined evidence" database using a search strategy focused on answering clinical questions relevant to ABS and employing a combination of Boolean searching on human-indexed thesaurus

CCHMC Grading Scale			
M	Meta-analysis or Systematic Review	S	Review Article
A	Randomized controlled trial: large sample	E	Expert opinion or consensus
B	Randomized controlled trial: small sample	F	Basic Laboratory Research
C	Prospective trial or large case series	L	Legal requirement
D	Retrospective analysis	Q	Decision analysis
O	Other evidence	X	No evidence

terms (MeSH headings using an OVID Medline interface) and "natural language" searching on words in the title, abstract, and indexing terms. The citations were reduced by: eliminating duplicates, review articles, non-English articles, and adult articles. The resulting abstracts were reviewed by a methodologist to eliminate low quality and irrelevant citations. During the course of the guideline development, additional clinical questions were generated and subjected to the search process, and some relevant review articles were identified. April, 2001 was the last date for which literature was reviewed for the previous version of this guideline. The details of previous review strategies are not documented. However, all original citations were reviewed for appropriateness to this revision.

Appropriate companion documents have been developed to assist in the effective dissemination and implementation of the guideline. Experience with implementation of the original publication of this guideline has provided learnings which have been incorporated into this revision. The outcome measures monitored as of the revision publication date are:

- percent of guideline-eligible patients seen in the Emergency Department who are prescribed antibiotics who have had symptoms ≥ 10 days duration or who are severely ill;
- percent of guideline-eligible patients seen in the Emergency Department and with symptoms ≥ 10 days duration who are prescribed either high-dose amoxicillin or high-dose amoxicillin-clavulanate.

Once the guideline has been in place for three years, the development team reconvenes to explore the continued validity of the guideline. This phase can be initiated at any point that evidence indicates a critical change is needed.

Recommendations have been formulated by a consensus process directed by best evidence, patient and family preference and clinical expertise. During formulation of these guidelines, the team members have remained cognizant of controversies and disagreements over the management of these patients. They have tried to resolve controversial issues by consensus where possible and, when not possible, to offer optional approaches to care in the form of information that includes best supporting evidence of efficacy for alternative choices.

The guidelines have been reviewed and approved by clinical experts not involved in the development process, and other individuals as appropriate to their intended purposes.

The guideline was developed without external funding. All Team Members and Clinical Effectiveness support staff listed have

declared whether they have any conflict of interest and none were identified.

Copies of this Evidence-Based Care Guideline (EBCG) and its companion documents are available online and may be distributed by any organization for the global purpose of improving child health outcomes. Website address:

<http://www.cincinnatichildrens.org/svc/alpha/h/health-policy/ev-based/default.htm> . Examples of approved uses of the EBCG include the following:

- copies may be provided to anyone involved in the organization's process for developing and implementing evidence-based care guidelines;
- hyperlinks to the CCHMC website may be placed on the organization's website;
- the EBCG may be adopted or adapted for use within the organization, provided that CCHMC receives appropriate attribution on all written or electronic documents; and
- copies may be provided to patients and the clinicians who manage their care.

Notification of CCHMC at HPCEInfo@cchmc.org for any EBCG, or its companion documents, adopted, adapted, implemented or hyperlinked by the organization is appreciated.

NOTE: These recommendations result from review of literature and practices current at the time of their formulations. This guideline does not preclude using care modalities proven efficacious in studies published subsequent to the current revision of this document. This document is not intended to impose standards of care preventing selective variances from the guidelines to meet the specific and unique requirements of individual patients. Adherence to this guideline is voluntary. The physician in light of the individual circumstances presented by the patient must make the ultimate judgment regarding the priority of any specific procedure.

For more information about these guidelines, their supporting evidences and the guideline development process, contact the Health Policy & Clinical Effectiveness office at: 513-636-2501 or HPCEInfo@cchmc.org

REFERENCES

Note: When using the electronic version of this document, “_____” refers to journal articles that have a hyperlink to the abstract.

- AAP:** American Academy of Pediatrics. Committee on Quality Improvement; clinical practice guideline: management of sinusitis. *Pediatrics*, 108(3): 798-808, 2001, [S] _____.
- AAP:** Use of codeine- and dextromethorphan-containing cough remedies in children. American Academy of Pediatrics. Committee on Drugs. *Pediatrics*, 99(6): 918-20, 1997, [S,E] _____.
- Aitken, M., and Taylor, J. A.:** Prevalence of clinical sinusitis in young children followed up by primary care pediatricians. *Arch Pediatr Adolesc Med*, 152(3): 244-8, 1998, [C] _____.
- Anon, J. B.; Jacobs, M. R.; Poole, M. D.; Ambrose, P. G.; Benninger, M. S.; Hadley, J. A.; and Craig, W. A.:** Antimicrobial treatment guidelines for acute bacterial rhinosinusitis. *Otolaryngol Head Neck Surg*, 130(1 Suppl): 1-45, 2004, [S,E] _____.
- Barden, L. S.; Dowell, S. F.; Schwartz, B.; and Lackey, C.:** Current attitudes regarding use of antimicrobial agents: results from physician's and parents' focus group discussions. *Clin Pediatr (Phila)*, 37(11): 665-71, 1998, [C] _____.
- Bernard, D. W.; Goepf, J. G.; Duggan, A. K.; Serwint, J. R.; and Rowe, P. C.:** Is oral albuterol effective for acute cough in non-asthmatic children? *Acta Paediatr*, 88(4): 465-7, 1999, [B] _____.
- Breiman, R. F.; Butler, J. C.; Tenover, F. C.; Elliott, J. A.; and Facklam, R. R.:** Emergence of drug-resistant pneumococcal infections in the United States. *JAMA*, 271(23): 1831-5, 1994, [C] _____.
- Brook, I.:** Bacteriologic features of chronic sinusitis in children. *JAMA*, 246(9): 967-9, 1981, [C] _____.
- Brook, I.; Gooch, W. M.; Jenkins, S. G.; Pichichero, M. E.; Reiner, S. A.; Sher, L.; and Yamauchi, T.:** Medical management of acute bacterial sinusitis. Recommendations of a clinical advisory committee on pediatric and adult sinusitis. *Ann Otol Rhinol Laryngol Suppl*, 182: 2-20, 2000, [S,E] _____.
- Butler, J. C.; Hofmann, J.; Cetron, M. S.; Elliott, J. A.; Facklam, R. R.; and Breiman, R. F.:** The continued emergence of drug-resistant *Streptococcus pneumoniae* in the United States: an update from the Centers for Disease Control and Prevention's Pneumococcal Sentinel Surveillance System. *J Infect Dis*, 174(5): 986-93, 1996, [C] _____.
- Chang, A. B.; Phelan, P. D.; Carlin, J. B.; Sawyer, S. M.; and Robertson, C. F.:** A randomised, placebo controlled trial of inhaled salbutamol and beclomethasone for recurrent cough. *Arch Dis Child*, 79(1): 6-11, 1998, [B] _____.
- Cincinnati Children's Hospital Medical Center:** Antibiotic Susceptibility Report. 2005, [O] _____.
- Clement, P. A. et al.:** Management of rhinosinusitis in children: consensus meeting, Brussels, Belgium, September 13, 1996. *Arch Otolaryngol Head Neck Surg*, 124(1): 31-4, 1998, [E] _____.
- Contopoulos-Ioannidis, D. G.; Ioannidis, J. P.; and Lau, J.:** Acute sinusitis in children: current treatment strategies. *Paediatr Drugs*, 5(2): 71-80, 2003, [S] _____.
- Dagan, R.; Hoberman, A.; Johnson, C.; Leibovitz, E. L.; Arguedas, A.; Rose, F. V.; Wynne, B. R.; and Jacobs, M. R.:** Bacteriologic and clinical efficacy of high dose amoxicillin/clavulanate in children with acute otitis media. *Pediatr Infect Dis J*, 20(9): 829-37, 2001, [C] _____.
- Dagan, R.; Johnson, C. E.; McLinn, S.; Abughali, N.; Feris, J.; Leibovitz, E.; Burch, D. J.; and Jacobs, M. R.:** Bacteriologic and clinical efficacy of amoxicillin/clavulanate vs. azithromycin in acute otitis media. *Pediatr Infect Dis J*, 19(2): 95-104, 2000, [A] _____.
- Davies, M. J.; Fuller, P.; Picciotto, A.; and McKenzie, S. A.:** Persistent nocturnal cough: randomised controlled trial of high dose inhaled corticosteroid. *Arch Dis Child*, 81(1): 38-44, 1999, [B] _____.
- Diament, M. J.:** The diagnosis of sinusitis in infants and children: x-ray, computed tomography, and magnetic resonance imaging. Diagnostic imaging of pediatric sinusitis. *J Allergy Clin Immunol*, 90(3 Pt 2): 442-4, 1992, [E] _____.
- Diament, M. J.; Senac, M. O.; Gilsanz, V.; Baker, S.; Gillespie, T.; and Larsson, S.:** Prevalence of incidental paranasal sinuses opacification in pediatric patients: a CT study. *J Comput Assist Tomogr*, 11(3): 426-31, 1987, [C] _____.
- Dowell, S. F.; Butler, J. C.; Giebink, G. S.; Jacobs, M. R.; Jernigan, D.; Musher, D. M.; Rakowsky, A.; and Schwartz, B.:** Acute otitis media: management and surveillance in an era of pneumococcal resistance--a report from the Drug-resistant *Streptococcus pneumoniae* Therapeutic Working Group. *Pediatr Infect Dis J*, 18(1): 1-9, 1999, [E] _____.
- Dowell, S. F.; Schwartz, B.; and Phillips, W. R.:** Appropriate use of antibiotics for URIs in children: Part I. Otitis media and acute sinusitis. The Pediatric URI Consensus Team. *Am Fam Physician*, 58(5): 1113-8, 1123, 1998a, [E] _____.
- Dowell, S. F.; Schwartz, B.; and Phillips, W. R.:** Appropriate use of antibiotics for URIs in children: Part II. Cough, pharyngitis and the common cold. The Pediatric URI Consensus Team. *Am Fam Physician*, 58(6): 1335-42, 1345, 1998b, [E] _____.
- Engels, E. A.; Terrin, N.; Barza, M.; and Lau, J.:** Meta-analysis of diagnostic tests for acute sinusitis. *Journal of Clinical Epidemiology*, 53(8): 852-62, 2000, [M] _____.
- Fearon, B.; Edmonds, B.; and Bird, R.:** Orbital-facial complications of sinusitis in children. *Laryngoscope*, 89(6 Pt 1): 947-53, 1979, [S] _____.
- Felmingham, D.; White, A. R.; Jacobs, M. R.; Appelbaum, P. C.; Poupard, J.; Miller, L. A.; and Gruneberg, R. N.:** The Alexander Project: the benefits from a decade of surveillance. *J Antimicrob Chemother*, 56 Suppl 2: ii3-ii21, 2005, [D] _____.

26. **Fireman, P.:** Diagnosis of sinusitis in children: emphasis on the history and physical examination. *J Allergy Clin Immunol*, 90(3 Pt 2): 433-6, 1992, [S] _____.
27. **Friedland, I. R., and McCracken, G. H.:** Management of infections caused by antibiotic-resistant *Streptococcus pneumoniae*. *N Engl J Med*, 331(6): 377-82, 1994, [E] _____.
28. **Gadomski, A., and Horton, L.:** The need for rational therapeutics in the use of cough and cold medicine in infants. *Pediatrics*, 89(4 Pt 2): 774-6, 1992, [O] _____.
29. **Garbutt, J. M.; Gellman, E. F.; and Littenberg, B.:** The development and validation of an instrument to assess acute sinus disease in children. *Qual Life Res*, 8(3): 225-33, 1999, [C] _____.
30. **Garbutt, J. M.; Goldstein, M.; Gellman, E.; Shannon, W.; and Littenberg, B.:** A randomized, placebo-controlled trial of antimicrobial treatment for children with clinically diagnosed acute sinusitis. *Pediatrics*, 107(4): 619-25, 2001, [B] _____.
31. **Gay, K.; Baughman, W.; Miller, Y.; Jackson, D.; Whitney, C. G.; Schuchat, A.; Farley, M. M.; Tenover, F.; and Stephens, D. S.:** The emergence of *Streptococcus pneumoniae* resistant to macrolide antimicrobial agents: a 6-year population-based assessment. *J Infect Dis*, 182(5): 1417-24, 2000, [D] _____.
32. **Glasier, C. M.; Ascher, D. P.; and Williams, K. D.:** Incidental paranasal sinus abnormalities on CT of children: clinical correlation. *AJNR Am J Neuroradiol*, 7(5): 861-4, 1986, [C] _____.
33. **Glasier, C. M.; Mallory, G. B.; and Steele, R. W.:** Significance of opacification of the maxillary and ethmoid sinuses in infants. *J Pediatr*, 114(1): 45-50, 1989, [C] _____.
34. **Gross, G. W.; McGeady, S. J.; Kerut, T.; and Ehrlich, S. M.:** Limited-slice CT in the evaluation of paranasal sinus disease in children. *Am J Roentgenol*, 156(2): 367-9, 1991, [C] _____.
35. **Group Health Associates:** Personal communication. 2004, [O].
36. **Gungor, A., and Corey, J. P.:** Pediatric sinusitis: a literature review with emphasis on the role of allergy. *Otolaryngol Head Neck Surg*, 116(1): 4-15, 1997, [S] _____.
37. **Gwaltney, J. M.; Phillips, C. D.; Miller, R. D.; and Riker, D. K.:** Computed tomographic study of the common cold. *N Engl J Med*, 330(1): 25-30, 1994, [D] _____.
38. **Hamm, R. M.; Hicks, R. J.; and Bembem, D. A.:** Antibiotics and respiratory infections: are patients more satisfied when expectations are met? *J Fam Pract*, 43(1): 56-62, 1996, [C] _____.
39. **Herrmann, B. W., and Forsen, J. W., Jr.:** Simultaneous intracranial and orbital complications of acute rhinosinusitis in children. *International Journal of Pediatric Otorhinolaryngology*, 68(5): 619-25, 2004, [D] _____.
40. **Hnatuk, L. A.; Macdonald, R. E.; and Papsin, B. C.:** Isolated sphenoid sinusitis: the Toronto Hospital for Sick Children experience and review of the literature. *J Otolaryngol*, 23(1): 36-41, 1994, [S] _____.
41. **Ioannidis, J. P., and Lau, J.:** Technical report: evidence for the diagnosis and treatment of acute uncomplicated sinusitis in children: a systematic overview. *Pediatrics*, 108(3): E57, 2001, [M] _____.
42. **Isaacson, G.:** Sinusitis in childhood. *Pediatr Clin North Am*, 43(6): 1297-318, 1996, [S] _____.
43. **Jacobs, M. R.; Bajaksouzian, S.; Zilles, A.; Lin, G.; Pankuch, G. A.; and Appelbaum, P. C.:** Susceptibilities of *Streptococcus pneumoniae* and *Haemophilus influenzae* to 10 oral antimicrobial agents based on pharmacodynamic parameters: 1997 U.S. Surveillance study. *Antimicrob Agents Chemother*, 43(8): 1901-8, 1999, [D] _____.
44. **Jacobs, M. R.; Felmingham, D.; Appelbaum, P. C.; and Gruneberg, R. N.:** The Alexander Project 1998-2000: susceptibility of pathogens isolated from community-acquired respiratory tract infection to commonly used antimicrobial agents. *J Antimicrob Chemother*, 52(2): 229-46, 2003, [D] _____.
45. **Jannert, M.; Andreasson, L.; Helin, I.; and Pettersson, H.:** Acute sinusitis in children—symptoms, clinical findings and bacteriology related to initial radiologic appearance. *Int J Pediatr Otorhinolaryngol*, 4(2): 139-48, 1982, [C] _____.
46. **Kakish, K. S.; Mahafza, T.; Batiha, A.; Ekteish, F.; and Daoud, A.:** Clinical sinusitis in children attending primary care centers. *Pediatr Infect Dis J*, 19(11): 1071-4, 2000, [C] _____.
47. **Kernan, W. N.; Viscoli, C. M.; Brass, L. M.; Broderick, J. P.; Brott, T.; Feldmann, E.; Morgenstern, L. B.; Wilterdink, J. L.; and Horwitz, R. I.:** Phenylpropanolamine and the risk of hemorrhagic stroke. *N Engl J Med*, 343(25): 1826-32, 2000, [D] _____.
48. **Kogutt, M. S., and Swischuk, L. E.:** Diagnosis of sinusitis in infants and children. *Pediatrics*, 52(1): 121-4, 1973, [D] _____.
49. **Lau, J. et al.:** Diagnosis and treatment of acute bacterial rhinosinusitis. *Evid Rep Technol Assess (Summ)*, (9): 1-5, 1999, [S] _____.
50. **Leopold, D. A.:** Clinical course of acute maxillary sinusitis documented by sequential MRI scanning. *Am J Rhinol*, 8(1): 19-28, 1994, [C].
51. **Local Expert Consensus:** During guideline development timeframe. [E].
52. **Loughlin, J.; Poullos, N.; Napalkov, P.; Wegmuller, Y.; and Monto, A. S.:** A study of influenza and influenza-related complications among children in a large US health insurance plan database. *Pharmacoeconomics*, 21(4): 273-83, 2003, [D] _____.
53. **Lusk, R. P., and Stankiewicz, J. A.:** Pediatric rhinosinusitis. *Otolaryngol Head Neck Surg*, 117(3 Pt 2): S53-7, 1997, [S] _____.

54. **Macfarlane, J.; Holmes, W.; Macfarlane, R.; and Britten, N.:** Influence of patients' expectations on antibiotic management of acute lower respiratory tract illness in general practice: questionnaire study. *BMJ*, 315(7117): 1211-4, 1997, [C] _____.
55. **Mangione-Smith, R.; McGlynn, E. A.; Elliott, M. N.; Krogstad, P.; and Brook, R. H.:** The relationship between perceived parental expectations and pediatrician antimicrobial prescribing behavior. *Pediatrics*, 103(4 Pt 1): 711-8, 1999, [C] _____.
56. **Maresh, M. M.:** Paranasal Sinuses from Birth to Late Adolescence: II. Clinical and Roentgenographic Evidence of Infection. *American Journal of Diseases of Children*, 60: 841-861, 1940, [D].
57. **McAlister, W. H. et al.:** Sinusitis in the pediatric population. American College of Radiology. ACR Appropriateness Criteria. *Radiology*, 215 Suppl: 811-8, 2000, [E] _____.
58. **McCormick, D. P.; John, S. D.; Swischuk, L. E.; and Uchida, T.:** A double-blind, placebo-controlled trial of decongestant-antihistamine for the treatment of sinusitis in children. *Clin Pediatr (Phila)*, 35(9): 457-60, 1996, [B] _____.
59. **McLean, D. C.:** Sinusitis in children. Lessons from twenty-five patients. *Clin Pediatr (Phila)*, 9(6): 342-5, 1970, [D] _____.
60. **Morris, P., and Leach, A.:** Antibiotics for persistent nasal discharge (rhinosinusitis) in children. *Cochrane Database Syst Rev*: CD001094, 2005, [M] _____.
61. **Morton, J. L., and Schultz, A. A.:** Healthy Hands: Use of alcohol gel as an adjunct to handwashing in elementary school children. *J Sch Nurs*, 20(3): 161-7, 2004, [A] _____.
62. **Muntz, H. R., and Lusk, R. P.:** Bacteriology of the ethmoid bullae in children with chronic sinusitis. *Arch Otolaryngol Head Neck Surg*, 117(2): 179-81, 1991, [D] _____.
63. **Myer, C. M., 3rd; Herwig, S. O.; Seid, A. B.; Cotton, R. T.; and Towbin, R. B.:** Acute sphenoid sinusitis in children. *Am J Otolaryngol*, 3(3): 223-6, 1982, [S] _____.
64. **Nash, D., and Wald, E.:** Sinusitis. *Pediatr Rev*, 22(4): 111-7, 2001, [S] _____.
65. **Nelson, C. T.; Mason, E. O.; and Kaplan, S. L.:** Activity of oral antibiotics in middle ear and sinus infections caused by penicillin-resistant *Streptococcus pneumoniae*: implications for treatment. *Pediatr Infect Dis J*, 13(7): 585-9, 1994, [C] _____.
66. **Nuutinen, J.; Holopainen, E.; Haahtela, T.; Ruoppi, P.; and Silvasti, M.:** Balanced physiological saline in the treatment of chronic rhinitis. *Rhinology*, 24(4): 265-9, 1986, [C] _____.
67. **Nyquist, A. C.; Gonzales, R.; Steiner, J. F.; and Sande, M. A.:** Antibiotic prescribing for children with colds, upper respiratory tract infections, and bronchitis. *JAMA*, 279(11): 875-7, 1998, [D] _____.
68. **Odita, J. C.; Akamaguna, A. I.; Ogisi, F. O.; Amu, O. D.; and Ugbodaga, C. I.:** Pneumatisation of the maxillary sinus in normal and symptomatic children. *Pediatr Radiol*, 16(5): 365-7, 1986, [C] _____.
69. **Ott, N. L.; O'Connell, E. J.; Hoffmans, A. D.; Beatty, C. W.; and Sachs, M. I.:** Childhood sinusitis. *Mayo Clin Proc*, 66(12): 1238-47, 1991, [S] _____.
70. **Oxford, L. E., and McClay, J.:** Complications of acute sinusitis in children. *Otolaryngology - Head & Neck Surgery*, 133(1): 32-7, 2005, [D] _____.
71. **Parsons, D. S., and Wald, E. R.:** Otitis media and sinusitis: similar diseases. *Otolaryngol Clin North Am*, 29(1): 11-25, 1996, [E] _____.
72. **Paul, I. M.; Yoder, K. E.; Crowell, K. R.; Shaffer, M. L.; McMillan, H. S.; Carlson, L. C.; Dilworth, D. A.; and Berlin, C. M., Jr.:** Effect of dextromethorphan, diphenhydramine, and placebo on nocturnal cough and sleep quality for coughing children and their parents. *Pediatrics*, 114(1): e85-90, 2004, [A] _____.
73. **Pichichero, M. E.; McLinn, S.; Aronovitz, G.; Fiddes, R.; Blumer, J.; Nelson, K.; and Dashefsky, B.:** Cefprozil treatment of persistent and recurrent acute otitis media. *Pediatr Infect Dis J*, 16(5): 471-8, 1997, [C] _____.
74. **Pigret, D., and Jankowski, R.:** Management of post-ethmoidectomy crust formation: randomized single-blind clinical trial comparing pressurized seawater versus antiseptic/mucolytic saline. *Rhinology*, 34(1): 38-40, 1996, [B] _____.
75. **Postma, G. N.; Chole, R. A.; and Nemzek, W. R.:** Reversible blindness secondary to acute sphenoid sinusitis. *Otolaryngol Head Neck Surg*, 112(6): 742-6, 1995, [E] _____.
76. **Rak, K. M.; Newell, J. D., 2nd; Yakes, W. F.; Damiano, M. A.; and Luethke, J. M.:** Paranasal sinuses on MR images of the brain: significance of mucosal thickening. *Am J Roentgenol*, 156(2): 381-4, 1991, [C] _____.
77. **Roberts, C. R.; Imrey, P. B.; Turner, J. D.; Hosokawa, M. C.; and Alster, J. M.:** Reducing physician visits for colds through consumer education. *JAMA*, 250(15): 1986-9, 1983, [A] _____.
78. **Roberts, L.; Smith, W.; Jorm, L.; Patel, M.; Douglas, R. M.; and McGilchrist, C.:** Effect of infection control measures on the frequency of upper respiratory infection in child care: a randomized, controlled trial. *Pediatrics*, 105(4 Pt 1): 738-42, 2000, [A] _____.
79. **Ros, S. P.; Herman, B. E.; and Azar-Kia, B.:** Acute sinusitis in children: is the Water's view sufficient? *Pediatr Radiol*, 25(4): 306-7, 1995, [D] _____.
80. **Rosenfeld, E. A., and Rowley, A. H.:** Infectious intracranial complications of sinusitis, other than meningitis, in children: 12-year review. *Clin Infect Dis*, 18(5): 750-4, 1994, [D] _____.

81. **Schrag, S. J.; Pena, C.; Fernandez, J.; Sanchez, J.; Gomez, V.; Perez, E.; Feris, J. M.; and Besser, R. E.:** Effect of short-course, high-dose amoxicillin therapy on resistant pneumococcal carriage: a randomized trial. *JAMA*, 286(1): 49-56, 2001, [A] _____
82. **Schroeder, K., and Fahey, T.:** Over-the-counter medications for acute cough in children and adults in ambulatory settings. *Cochrane Database Syst Rev*, (4): CD001831, 2004, [M] _____
83. **Schwartz, R. H.; Pitkaranta, A.; and Winther, B.:** Computed tomography imaging of the maxillary and ethmoid sinuses in children with short-duration purulent rhinorrhea. *Otolaryngol Head Neck Surg*, 124(2): 160-3, 2001, [C] _____
84. **Sellers, S. L.; Goldberg, S.; and Seid, A. B.:** Transnasal sphenoidotomy. *S Afr Med J*, 49(33): 1351-5, 1975, [S] _____
85. **Shopfner, C. E., and Rossi, J. O.:** Roentgen evaluation of the paranasal sinuses in children. *Am J Roentgenol Radium Ther Nucl Med*, 118(1): 176-86, 1973, [C] _____
86. **Shoseyov, D.; Bibi, H.; Shai, P.; Shoseyov, N.; Shazberg, G.; and Hurvitz, H.:** Treatment with hypertonic saline versus normal saline nasal wash of pediatric chronic sinusitis. *J Allergy Clin Immunol*, 101(5): 602-5, 1998, [B] _____
87. **Steele, R. W.; Estrada, B.; Begue, R. E.; Mirza, A.; Travillion, D. A.; and Thomas, M. P.:** A double-blind taste comparison of pediatric antibiotic suspensions. *Clin Pediatr (Phila)*, 36(4): 193-9, 1997, [O] _____
88. **Taylor, J. A.; Novack, A. H.; Almquist, J. R.; and Rogers, J. E.:** Efficacy of cough suppressants in children. *J Pediatr*, 122(5 Pt 1): 799-802, 1993, [B] _____
89. **Trepka, M. J.; Belongia, E. A.; Chyou, P. H.; Davis, J. P.; and Schwartz, B.:** The effect of a community intervention trial on parental knowledge and awareness of antibiotic resistance and appropriate antibiotic use in children. *Pediatrics*, 107(1): E6, 2001, [C] _____
90. **Van Buchem, F. L.; Peeters, M. F.; and Knotnerus, J. A.:** Maxillary sinusitis in children. *Clin Otolaryngol Allied Sci*, 17(1): 49-53, 1992, [C] _____
91. **Varonen, H., and Sainio, S.:** Patients' and physicians' views on the management of acute maxillary sinusitis. *Scandinavian Journal of Primary Health Care*, 22(1): 22-6, 2004, [O] _____
92. **Vazquez, E.; Creixell, S.; Carreno, J. C.; Castellote, A.; Figueras, C.; Pumarola, F.; Poch, J. M.; and Lucaya, J.:** Complicated acute pediatric bacterial sinusitis: Imaging updated approach. *Current Problems in Diagnostic Radiology*, 33(3): 127-45, 2004, [D] _____
93. **Wald, E. R.:** Management of sinusitis in infants and children. *Pediatr Infect Dis J*, 7(6): 449-52, 1988, [E] _____
94. **Wald, E. R.:** Sinusitis in children. *Isr J Med Sci*, 30(5-6): 403-7, 1994, [S] _____
95. **Wald, E. R.; Chiponis, D.; and Ledesma-Medina, J.:** Comparative effectiveness of amoxicillin and amoxicillin-clavulanate potassium in acute paranasal sinus infections in children: a double-blind, placebo-controlled trial. *Pediatrics*, 77(6): 795-800, 1986, [B] _____
96. **Wald, E. R.; Guerra, N.; and Byers, C.:** Upper respiratory tract infections in young children: duration of and frequency of complications. *Pediatrics*, 87(2): 129-33, 1991, [C] _____
97. **Wald, E. R.; Milmo, G. J.; Bowen, A.; Ledesma-Medina, J.; Salamon, N.; and Bluestone, C. D.:** Acute maxillary sinusitis in children. *N Engl J Med*, 304(13): 749-54, 1981, [B] _____
98. **Wald, E. R.; Reilly, J. S.; Casselbrant, M.; Ledesma-Medina, J.; Milmo, G. J.; Bluestone, C. D.; and Chiponis, D.:** Treatment of acute maxillary sinusitis in childhood: a comparative study of amoxicillin and cefaclor. *J Pediatr*, 104(2): 297-302, 1984, [B] _____
99. **Wasserfallen, J. B.; Livio, F.; and Zanetti, G.:** Acute rhinosinusitis : a pharmaco-economic review of antibacterial use. *Pharmacoeconomics*, 22(13): 829-37, 2004, [S] _____
100. **Watt-Boolsen, S., and Karle, A.:** The clinical use of radiological examination of the maxillary sinuses. *Clin Otolaryngol Allied Sci*, 2(1): 41-3, 1977, [C] _____
101. **Whitney, C. G. et al.:** Increasing prevalence of multidrug-resistant *Streptococcus pneumoniae* in the United States. *N Engl J Med*, 343(26): 1917-24, 2000, [D] _____
102. **Williams, J. W., and Simel, D. L.:** Does this patient have sinusitis? Diagnosing acute sinusitis by history and physical examination. *JAMA*, 270(10): 1242-6, 1993, [S] _____
103. **Wippold, F. J.; Levitt, R. G.; Evens, R. G.; Korenblat, P. E.; Hodges, F. J.; and Jost, R. G.:** Limited coronal CT: an alternative screening examination for sinonasal inflammatory disease. *Allergy Proc*, 16(4): 165-9, 1995, [C] _____
104. **Wolf, G.; Anderhuber, W.; and Kuhn, F.:** Development of the paranasal sinuses in children: implications for paranasal sinus surgery. *Ann Otol Rhinol Laryngol*, 102(9): 705-11, 1993, [F] _____
105. **Wu, J. H.; Howard, D. H.; McGowan, J. E., Jr.; Frau, L. M.; and Dai, W. S.:** Patterns of health care resource utilization after macrolide treatment failure: results from a large, population-based cohort with acute sinusitis, acute bronchitis, and community-acquired pneumonia. *Clinical Therapeutics*, 26(12): 2153-62, 2004, [D] _____
106. **Wyllie, J. W., 3rd; Kern, E. B.; and Djalilian, M.:** Isolated sphenoid sinus lesions. *Laryngoscope*, 83(8): 1252-65, 1973, [S] _____
107. **Zacharisen, M. C., and Kelly, K. J.:** Allergic and infectious pediatric sinusitis. *Pediatr Ann*, 27(11): 759-66, 1998, [S] _____